

Amar Ojha<sup>1,2,3</sup>; Teague Henry<sup>4,5</sup>; Rasim Diler<sup>3</sup>; Cecile D. Ladouceur<sup>1,2,3,6</sup>

1. Center for Neuroscience; 2. Center for Neural Basis of Cognition; 3. Department of Psychiatry, University of Pittsburgh; 4. Department of Psychology, University of Virginia; 5. School of Data Science, University of Virginia; 6. Department of Psychology, University of Pittsburgh

## BACKGROUND

- Anhedonia—the reduced capacity for pleasure—is a common<sup>1</sup> and debilitating<sup>2</sup> feature of adolescent depression.
- Previous work has implicated altered reward circuitry<sup>3</sup>, which undergoes protracted maturation through adolescence<sup>4</sup>, in anhedonia; however, the role of striatal neurophysiology in remains unclear.
- Measures derived from resting-state functional magnetic resonance imaging (rsfMRI) can non-invasively provide *in vivo* measurements of various aspects of striatal neurophysiology.

**Hypotheses:** (1) Lower striatal regional homogeneity (ReHo) (Fig. 1, path b) and lower dopamine function (T2\*) intensity (Fig. 1, path c) will be associated with higher levels of anhedonia. (2) Striatal tissue iron will moderate this relationship, such that lower ReHo will be most strongly linked to high levels of anhedonia at low level of tissue iron.

## METHODS

- Sample: 75 adolescents (M age = 15.30 (1.50), 46 F) participated in the study, of whom 56 scored  $\geq 40$  on the Children's Depression Rating Scale-Revised (CDRS-R)<sup>5</sup> and 19 reported no current/past self/parent psychiatric diagnosis.
- Anhedonia was assessed using the Snaith-Hamilton Pleasure Scale (SHAPS)<sup>6</sup> and other depressive symptoms using the Mood and Feelings Questionnaire (MFQ)<sup>7</sup>.
- We derived two measures from two 6-minute resting-state functional magnetic resonance imaging (rsfMRI) scans:
  - ReHo to assess regional voxel synchronization<sup>8</sup>
  - Normalized T2\* signal to assess tissue iron<sup>9</sup>
- Data were preprocessed using *fmriprep*<sup>10</sup>.
- We used a voxel-wise moderated mediation approach to examine the relationships between T2\* intensity, ReHo, and anhedonia severity, controlling for age, sex, and depressive symptoms besides anhedonia (Fig. 1).
- Clusterwise permutation testing accounted for multiple comparison corrections.




Figure 1. Path diagram for moderated mediation model.

## RESULTS

- Reduced striatal tissue iron in the left putamen was associated with higher levels of anhedonia (Fig. 2).
- Reduced ReHo was associated with higher levels of anhedonia in adolescents with higher striatal tissue iron in the right caudate, and with lower levels of anhedonia in adolescents with lower levels of tissue iron.




Figure 2. Simple slopes for T2\* intensity effect on SHAPS score (Hypothesis 1)




Figure 3. Simple slopes for ReHo effect on SHAPS score (Hypotheses 1 + 2)




Figure 4. Moderating effect of T2\* intensity on ReHo-SHAPS score simple slopes (Hypothesis 2)

## CONCLUSIONS

- Both lower striatal ReHo and striatal tissue iron were associated with anhedonia beyond other depressive symptoms.
- However, these effects were specific to the dorsal striatum and the direction of this relationship was contingent upon levels of striatal tissue iron.
- Future research is needed to determine the effectiveness of dopamine-targeted pharmacotherapy for adolescents with anhedonia and particularly those with altered dopaminergic functioning.

## REFERENCES

- Yorbik, O., Birmaher, B., Axelson, D., Williamson, D. E., & Ryan, N. D. (2004). Clinical characteristics of depressive symptoms in children and adolescents with major depressive disorder. *Journal of Clinical Psychiatry*, 65 (12), 1654-1659.
- Ely, B. A., Nguyen, T. N., Tobe, R. H., Walker, A. M., & Gabbay, V. (2021). Multimodal investigations of reward circuitry and anhedonia in adolescent depression. *Frontiers in Psychiatry*, 12, 678709.
- Forbes, E. E., Hariri, A. R., Martin, S. L., Silk, J. S., Moyses, D. L., Fisher, P. M., ... & Dahl, R. E. (2009). Altered striatal activation predicting real-world positive affect in adolescent major depressive disorder. *American Journal of Psychiatry*, 166 (1), 64-73.
- Larsen, B., & Luna, B. (2015). In vivo evidence of neurophysiological maturation of the human adolescent striatum. *Developmental Cognitive Neuroscience*, 12, 74-85.
- Poznanski, E. O., & Mokros, H. B. (1996). Children's depression rating scale, revised (CDRS-R)
- Snaith, R. P., Hamilton, M., Morley, S., Humayan, A., Hargreaves, D., & Trigwell, P. (1995). A scale for the assessment of hedonic tone the Snaith-Hamilton Pleasure Scale. *The British Journal of Psychiatry*, 167 (1), 99-103.
- Angold, A., & Costello, E. J. (1987). Mood and feelings questionnaire (MFQ). *Durham: Developmental Epidemiology Program, Duke University*.
- Zang, Y., Jiang, T., Lu, Y., He, Y., & Tian, L. (2004). Regional homogeneity approach to fMRI data analysis. *NeuroImage*, 22(1), 394-400.
- Haacke, E. M., Cheng, N. Y., House, M. J., Liu, Q., Neelavalli, J., Ogg, R. J., ... & Obenaus, A. (2005). Imaging iron stores in the brain using magnetic resonance imaging. *Magnetic Resonance Imaging*, 23(1), 1-25.
- Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., ... & Gorgolewski, K. J. (2019). fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature Methods*, 16 (1), 111-116.

## FUNDING

Research funded by NIMH R01MH111600 (mPIs: Ladouceur, Diler)

## ACKNOWLEDGEMENTS

The authors would like to thank the study participants and the staff of the CAN-D Lab who made this research possible.

